

Annihilation catastrophe: new critical phenomenon in diffusion-controlled dynamics

Boris M. Shipilevsky

Institute of Solid State Physics, Chernogolovka, Moscow district, 142432, Russia

For the last decade the reaction-diffusion system $A + B \rightarrow 0$, where unlike species A and B diffuse and irreversibly react in the bulk of a d -dimensional substrate, has acquired the status of one of the most popular objects in nonequilibrium statistical physics [1]. Two problems have been investigated most intensively: (i) dynamics of the $A + B \rightarrow 0$ annihilation in an infinite system with initially homogeneously (randomly) and equimolarly distributed reactants, in which below the critical dimension $d_c = 4$ a dynamical clustering develops (fluctuation-induced like-particle domains formation) and, as a result, an anomalous reaction deceleration arises (Ovchinnikov and Zeldovich, 1978 [2], Toussaint and Wilczek, 1983 [3]); (ii) behaviour of the dynamic reaction front in an infinite system with initially spatially separated reactants (Galfi and Racz, 1988 [4]), and structure of the steady state front in a finite system, at the ends of which are injected equal currents of A 's and B 's particles (Ben-Naim and Redner, 1992 [5], Cornell and Droz, 1993 [6]).

In 1994 it was discovered (Shipilevsky, 1994 [7]) that in another wide class of RD systems, where reaction and diffusion are spatially separated (i.e. reaction proceeds on the surface of the medium and diffusion proceeds in its bulk) the interplay between reaction and diffusion acquires qualitatively new features and leads to the threshold self-organizing dynamics of the $A + B \rightarrow 0$. It has been found that once particles A and B diffuse at different mobilities from the bulk of finite medium onto the surface and die on it by the reaction $A + B \rightarrow 0$, there should exist some threshold difference in the initial numbers of A and B particles, Δ_c , above which the loop of positive feedback is "switched on" and the process of their death, instead of usual deceleration, starts to *accelerate autocatalytically*. Recently, I have shown [8] that deceleration-acceleration transition is only a prelude to much more nontrivial dynamical effects, which take place far beyond the acceleration threshold. It has been demonstrated that in the diffusion-controlled limit $\Delta \rightarrow \infty$ a new critical phenomenon develops, which arises as a result of self-organizing explosive growth (drop) of the surface concentrations of, respectively, slow and fast particles (*concentration explosion*) and manifests itself in the form of an abrupt singular jump in the desorption flux relaxation rate. The aim of this report is to give an insight into the key features of this remarkable phenomenon which I call *annihilation catastrophe*.

Model. I consider a model, in which species A and B are supposed to be initially uniformly distributed in the bulk of infinitely extended slab of thickness 2ℓ . Both species diffuse to the surface ($X = \pm\ell$) and desorb as a result of surface reaction $A_{ads} + B_{ads} \rightarrow AB \rightarrow 0$. The rate of reaction is proportional to the product of surface concentrations $I = k\rho_A\rho_B = \kappa c_{As}c_{Bs}$, where the rates of surface-subsurface exchange are supposed to be rather high, so that quasiequilibrium $\rho_i = f_i c_{is}$ is always sustained. The system is

effectively one dimensional and the boundary conditions are determined from the equality of diffusion I^D and desorption I flux densities at the surface $I^D|_s = I$, i.e. it is assumed that the surface layer capacity can be neglected. Thus, introducing index "H" (*heavy*) for slower diffusing species and index "L" (*light*) for a faster one, the problem reads (by symmetry I consider the interval $[0, \ell]$ only)

$$\partial h / \partial \tau = \nabla^2 h \quad , \quad \partial l / \partial \tau = (1/p) \nabla^2 l, \quad (1)$$

$$\nabla h|_s = (1/p) \nabla l|_s = -h_s l_s, \quad (2)$$

with $\nabla(h, l)|_{x=0} = 0$ and the initial conditions $h(x, 0) = h_0$ and $l(x, 0) = l_0$. Here $h(x, \tau) = c_H/c_*$ and $l(x, \tau) = c_L/c_*$ are the reduced concentrations, $\nabla \equiv \partial/\partial x$, $x = X/\ell \in [0, 1]$ is the dimensionless coordinate, $\tau = D_H t/\ell^2$ is the dimensionless time, $p = D_H/D_L \leq 1$ is the ratio of species diffusivities, and $c_* = D_H/\kappa\ell$ is the characteristic concentration scale, from which diffusion starts to play an essential role. Boundary conditions (2) reflect the fact that particles disappear in pairs only, i.e. $J = h_s l_s = -\langle \dot{h} \rangle = -\langle \dot{l} \rangle$, where $J = I/I_*$ is the reduced desorption flux density and $I_* = \kappa c_*^2$ is its characteristic scale, therefore $\langle h \rangle - \langle l \rangle = \Delta = \text{const.}$, i.e., the excess amount stays "inert" in the bulk (here $\langle h \rangle = \int_0^1 h dx = \mathcal{N}_H/\mathcal{N}_*$ and $\langle l \rangle = \int_0^1 l dx = \mathcal{N}_L/\mathcal{N}_*$ are the total reduced numbers of particles in the bulk per unit of surface and $\mathcal{N}_* = c_* \ell = D_H/\kappa$ is the characteristic particle number scale). I consider here the annihilation dynamics for $\Delta > 0$ when the pair number N is dictated by the L particles number $N(\tau) = \mathcal{N}_{pair}/\mathcal{N}_* = N_L(\tau)$ so that in the final state all L particles disappear $N(\infty) = l(x, \infty) = 0$ and H particles are distributed uniformly with concentration $h(x, \infty) = \Delta$. I am interested first and foremost in the diffusion-controlled limit $N_0 \rightarrow \infty$ (note that diffusion-controlled limit $N_0 = l_0 = c_L(0)/c_* \rightarrow \infty$ means $c_*^{-1} = \kappa\ell/D_H \rightarrow \infty$). Moreover, I focus on the limit $n_0 = N_0/\Delta \gg 1$ when the initial numbers of H and L particles are almost equal, $r = h_0/l_0 = (1 + n_0)/n_0 \approx 1$. Precisely in this limit the particles interplay is most "intensive" and gives rise to the explosive dynamics, that is my prime interest here.

Transition point and self-accelerating annihilation. Assuming $N_0 \rightarrow \infty$ and $\mathcal{R} = r\sqrt{p} < 1$, I show [10], that for $n_0 \gg 1$ and $\gamma = 1 - \mathcal{R} \gg 1/\sqrt{N_0}$ at times $\tau \sim \tau_h = 1/N_0^2 \rightarrow 0$ the annihilation is followed by a sharp drop of h_s , crossing over to the H -diffusion-controlled regime, where the desorption flux density is governed by the rate of H particles diffusion to the system's surface. In the range $\tau_h \ll \tau \ll 1$, when diffusion of H particles proceeds in a semi-infinite medium regime, $J \approx h_0/\sqrt{\pi\tau} \approx J_0/\sqrt{\pi}(\tau_h/\tau)^{1/2}$ and for surface concentrations I find

$$h_s = (r/\gamma\sqrt{\pi\tau})(1 + \varrho + \dots), \quad l_s = l_0\gamma(1 - \varrho + \dots),$$

where $\varrho = 2r\sqrt{\tau/\pi}$ at $p \ll 1$. At $\tau > 1$ the slow power-law stage crosses over to the exponential relaxation stage $J = \mathcal{A}e^{-\omega_0\tau}(1 + e^{-8\omega_0\tau} + \dots)$ with amplitude $\mathcal{A} = 2h_0$, and I show that in the limit $\mathcal{A} \rightarrow \infty$ the surface concentrations achieve universal (Δ -*independent*) asymptotics

$$h_s = \Delta_c(1 + \phi + e^{-8\omega_0\tau} + \dots), \quad l_s = (\mathcal{A}/\Delta_c)e^{-\omega_0\tau}(1 - \phi + \dots) \quad (3)$$

where $|\phi|$ decays as $O(e^{-8\omega_0\tau})$ in the range $p \leq p_c = 4/9$ and as $O(e^{-(\pi^2/p-\omega_0)\tau})$ in the range $p_c < p < 1$ [10], and Δ_c obeys the expression [7,9]

$$\Delta_c = \sqrt{\omega_0/p} \tan(\sqrt{\omega_0 p})$$

($\omega_0 = \pi^2/4$ being the main eigenfrequency of the diffusion field relaxation under absorbing boundary conditions). This important result implies that in the diffusion-controlled limit $\mathcal{A} \rightarrow \infty$ the critical asymptotics $h_s = h_s^c = \Delta_c$ appears at *any* Δ as a *precursor* of long-time asymptotics $h_s|_{\tau \rightarrow \infty} = \Delta$, and precisely this intermediate asymptotics h_s^c *selects* the critical value $\Delta = \Delta_c = h_s^c$ below and above of which the relaxation dynamics of h_s is qualitatively different: at $\Delta < \Delta_c$ the surface concentration, h_s , escaping from the asymptotics h_s^c , decreases, reaching its limiting value Δ from *above*, whereas at $\Delta > \Delta_c$ the surface concentration, h_s , escaping from the asymptotics h_s^c , increases, reaching its limiting value Δ from *below* [7]. With neglect of rapidly decaying terms of $O(e^{-8\omega_0\tau}, e^{(\pi^2/p-\omega_0)\tau})$ the exact asymptotic expansion in power of $1/\mathcal{A}$ has the form [10]

$$h_s = \Delta_c(1 + \lambda + \dots), \quad l_s = (\mathcal{A}/\Delta_c)e^{-\omega_0\tau}(1 - \lambda + \dots), \quad N = (\mathcal{A}/\omega_0)e^{-\omega_0\tau}(1 - \mu\lambda + \dots),$$

$$J = \mathcal{A}e^{-\omega_0\tau}(1 - \sqrt{\omega_0} \tanh \sqrt{\omega_0}(\Delta - \Delta_c)^{-1}\lambda^2 + \dots), \quad (4)$$

with $\mu = \omega_0/\Delta_c \approx 1-p$ and $\lambda = \Delta_c(\Delta - \Delta_c)\mathcal{A}^{-1}e^{\omega_0\tau}$. These expressions and more detailed analysis in Ref.[7,9] suggest that at the crossing of the critical point $\Delta = \Delta_c$ not only the sign of $\dot{h}_s$ changes, but a growth of h_s (and with it the whole annihilation process) begins to accelerate *autocatalytically*, $\dot{h}_s = \omega_0(h_s - \Delta_c) + \dots$, as a result of "switching-on" of the loop of positive feedback: diffusion-induced growth of h_s accelerates the drop of l_s , the drop of l_s accelerates the growth of h_s and so on (*self-accelerating pairs death*). This phenomenon, interpreted in [7] as a new type of self-organization, develops the brighter the stronger Δ exceeds Δ_c , and in the limit $\Delta \rightarrow \infty$ it results in the explosive dynamics - *annihilation catastrophe* [8].

Annihilation catastrophe. Scaling and universality. According to (4) in the limit $\Delta \rightarrow \infty$ at the self-acceleration stage the rates of growth $\Omega_{Hs} = +d \ln h_s/d\tau$ and relaxation $\Omega_{Ls} = -d \ln l_s/d\tau$ of surface concentrations are unambiguously defined by the current value of the reduced pair number $n(\tau) = N(\tau)/\Delta$ and are mutually "compensated" so that with an accuracy to vanishingly small terms $O(1/n^2\Delta) \rightarrow 0$ the desorption flux relaxation rate $\tau_J^{-1} = -d \ln J/d\tau = \Omega_{Ls} - \Omega_{Hs} = \omega_0 + \dots$ is sustained constant. The central result of [8] is my finding, that at $\Delta \rightarrow \infty$ the relaxation rate τ_J^{-1} is sustained constant up to the time moment τ_* of reaching some critical pair number $n_*(p)$, in whose vicinity $\delta\tau \propto \Delta^{-2/5} \rightarrow 0$ the τ_J^{-1} abruptly jumps from $\tau_J^{-1} = \omega_0$ to $\tau_J^{-1} \rightarrow \infty$. I consider here the two consecutive stages of this phenomenon.

i) *Concentration explosion.* Following the work [8], I first show that condition $\tau_J^{-1} = \omega_0$ in the limit $\Delta \rightarrow \infty$ leads automatically to the arising of a *finite-time singularity*. Taking $J = \mathcal{A}e^{-\omega_0\tau}$ and satisfying (1),(2) to the leading order in Δ , I find

$$j = J/\Delta = -\dot{n} = \omega_0(1 + n), \quad h_s = \Delta_c(1 + n)/(n - n_*), \quad l_s = \mu\Delta(n - n_*), \quad (5)$$

where $n_* = (1 - \mu)/\mu \approx p/(1 - p)$. It is seen that h_s diverges as $n \rightarrow n_*$ and at the point of singularity exactly $j_* = \Delta_c$. From equality $J_* = 2h_0e^{-\omega_0\tau_*} = \Delta\Delta_c$ in accord with the numerical data of [10] we obtain

$$\tau_\star = (4/\pi^2) \ln[2(1+n_0)/\Delta_c].$$

Taking the origin of time at the point $\tau_\star$, i.e. introducing a relative time $\mathcal{T} = \tau - \tau_\star$, from (5) we come to remarkably simple laws $n = (1/\mu)e^{\omega_0|\mathcal{T}|} - 1$, $j = \Delta_c e^{\omega_0|\mathcal{T}|}$, $h_s = \Delta_c/(1 - e^{-\omega_0|\mathcal{T}|})$, $l_s = \Delta(e^{\omega_0|\mathcal{T}|} - 1)$, and, hence, $\Omega_{Hs} = \mu(h_s - \Delta_c) = \omega_0/(e^{\omega_0|\mathcal{T}|} - 1)$, whence it follows that for any $p < 1$ at the "distance" from the singularity point $|\mathcal{T}| \sim \omega_0^{-1}$ the dynamics of growth of h_s changes drastically: the initial exponential increase in the growth rate $\Omega_{Hs} = \omega_0 e^{-\omega_0|\mathcal{T}|}$ ($\Omega_{Ls} \approx \omega_0$) turns at $\omega_0|\mathcal{T}| \ll 1$ to a singular one

$$\Omega_{Hs} = \Omega_{Ls} = \mu h_s = 1/|\mathcal{T}|, \quad |\mathcal{T}| \rightarrow 0, \quad (6)$$

i.e., *concentration explosion* develops in the system. The striking feature of this phenomenon is that at times $\omega_0|\mathcal{T}| \ll 1$ the desorption flux practically reaches its critical value $J_\star$, therefore the concentration explosion develops, in fact, at a "frozen" flux $h_s l_s = J_\star$, $h_s = 1/\mu|\mathcal{T}|$, $l_s = \mu J_\star |\mathcal{T}|$, the consequence of which is *synchronization* of the growth and relaxation rates $\Omega_{Hs} = \Omega_{Ls} = \Omega_s$ during the explosion. A singular growth of h_s at the background of the diffusion flux $J_\star$, "frozen" on the explosion time scale, gives rise to a singular growth of the diffusion "antiflux", the calculation of which yields [8] $\delta J/J_\star \sim -1/\Delta|\mathcal{T}|^{3/2}$. As a result, in the desorption flux relaxation rate $\tau_J^{-1} = \omega_0 + [\tau_J^{-1}]$ the singularly growing term arises

$$[\tau_J^{-1}] \sim 1/\Delta|\mathcal{T}|^{5/2}, \quad (7)$$

whence it follows that the scenario described above holds down to $|\mathcal{T}|_{cat} \sim \Delta^{-2/5} \rightarrow 0$, beyond which a catastrophic growth of τ_J^{-1} begins.

ii) *Two universality classes of passing through the point of singularity.* Remarkably, the dynamics of passing through the point of singularity can be described analytically (which, in itself, is a rather rare event [11]) and it appears quite beautiful. I show analytically and confirm numerically that at the singularity point $\mathcal{T} = 0$ the rate of explosion Ω_{Hs} reaches a maximum Ω_{Hs}^M , the dynamics of passing through which proceeds in one of the two characteristic scaling regimes, depending on the value of parameter $\mathcal{K} = p^{3/2}\Delta/\Delta_c$.

1. *Annihilation catastrophe at "frozen" flux* ($\mathcal{K} \gg 1$). In the limit $\mathcal{K} \gg 1$ at times $|\mathcal{T}| \ll \mathcal{T}_L = p$ the particle number $n = n_\star(1 + \omega_0|\mathcal{T}|/p + \dots)$ is, in fact, "frozen" $n \approx n_\star$ and so in the process of explosion in the contracting as $\delta x_L \sim \sqrt{|\mathcal{T}|/p}$ nearsurface layer there arises a sharp gradient of L particles. I show that in this limit up to $\mathcal{T} \ll p$ the flux and the particle number are retained "frozen", $h_s l_s = J_\star$ and, hence, the explosion develops synchronously, $\Omega_{Hs} = \Omega_{Ls} = \Omega_s$, both *before* and *after* the singularity point. Equalizing the rates of change of H and L particles diffusion fluxes leads to the scaling

$$\Omega_s = \Omega_s^M S(\mathcal{T}/\mathcal{T}_f), \quad h_s/h_s^M = l_s^M/l_s = w(\mathcal{T}/\mathcal{T}_f), \quad (8)$$

where the scaling functions $S(\zeta) = 1/\sqrt{1+\zeta^2}$ and $w(\zeta) = \zeta + \sqrt{1+\zeta^2}$, the characteristic time $\mathcal{T}_f = 1/\Omega_s^M$, and the amplitudes Ω_s^M , h_s^M and l_s^M at the explosion maximum

$$h_s^M = (2/\mu)\Omega_s^M = p^{-1/4}\sqrt{\Delta\Delta_c}, \quad l_s^M = p^{1/4}\sqrt{\Delta\Delta_c}.$$

Two striking features of this result are symmetrical universalization of $\Omega_s = 1/|\mathcal{T}|$ beyond the scope of interval $[-\mathcal{T}_f, \mathcal{T}_f]$, $\mathcal{T}_f \rightarrow 0$ and remarkable symmetry $\mathcal{T} \leftrightarrow -\mathcal{T}$, $w \leftrightarrow 1/w$.

Singular "flash" of Ω_s leads to a singular jump in τ_J^{-1} , the dynamics of which is described by the scaling

$$[\tau_J^{-1}] = [\tau_J^{-1}]_* W(\mathcal{T}/\mathcal{T}_f), \quad [\tau_J^{-1}]_* = c_1(\Omega_s^M)^{5/2}/\Delta \propto \mu^{5/4} p^{-5/8} \Delta^{1/4}, \quad (9)$$

where in accord with the scaling function $W(\zeta) = c_2 \int_0^\infty d\theta / \sqrt{\theta} [1 + (\zeta - \theta)^2]^{3/2}$ the value of $[\tau_J^{-1}]$ reaches at $\mathcal{T} \sim \mathcal{T}_f$ a maximum $\max[\tau_J^{-1}] \sim [\tau_J^{-1}]_* \propto \Delta^{1/4} \rightarrow \infty$, and then it decays as $\sim 1/\sqrt{p\mathcal{T}}$. Essentially, that $\max[\tau_J^{-1}] \propto p^{-5/8}$, so the less is p the more brightly the effect is displayed.

2. Annihilation catastrophe at uniform L particles distribution. Flux breaking effect ($\mathcal{K} \ll 1$). If $\Delta \rightarrow \infty$ but $p \rightarrow 0$ so that $\mathcal{K} \ll 1$, the distribution of L species is retained uniform ($l_s = N, \Omega_{Ls} = h_s$) and the final stage of explosion is changed radically. I demonstrate that in this case an intermediate scaling takes place

$$\Omega_{Hs} = \Omega_{Hs}^M Q(\mathcal{T}/\mathcal{T}_*), \quad h_s = h_s^M v_H(\mathcal{T}/\mathcal{T}_*), \quad l_s = l_s^M v_L(\mathcal{T}/\mathcal{T}_*) \quad (10)$$

where the characteristic time $\mathcal{T}_* = 1/\Omega_{Hs}^M$ and amplitudes Ω_{Hs}^M, h_s^M and l_s^M at the explosion maximum

$$h_s^M = q\Omega_{Hs}^M = b_H \Delta^{2/3}, \quad l_s^M = b_L \Delta^{1/3},$$

with $q = 2.24, b_H = 1.63$ and $b_L = 0.73$. It is seen that in the limit $\mathcal{K} \ll 1$ the singular "flash" of Ω_{Hs} occurs much sharper, leading to more abrupt jump in τ_J^{-1}

$$[\tau_J^{-1}] = [\tau_J^{-1}]_* V(\mathcal{T}/\mathcal{T}_*), \quad [\tau_J^{-1}]_* = (q - 1)\Omega_{Hs}^M \propto \Delta^{2/3}, \quad (11)$$

which, in turn, gives rise to a qualitatively new phenomenon: the finite jump of the flux by a factor of ≈ 2 for the time $|\mathcal{T}|_{break} \sim \mathcal{T}_* \propto \Delta^{-2/3} \rightarrow 0$ (*flux breaking effect*).

As in the limit $\mathcal{K} \gg 1$ at $\mathcal{T} \gg p$ the relaxation rate τ_J^{-1} reaches its L -diffusion-controlled limit $\tau_J^{-1} \sim \omega_0/p$, and in the limit $\mathcal{K} \ll 1$ it always grows, I point out finally (and demonstrate numerically) that independently of $\mathcal{K}$ in the limit $p \rightarrow 0, \Delta \rightarrow \infty$ there arises a most spectacular consequence of the annihilation catastrophe: *an abrupt, practically instantaneous (on the scale of ω_0) disappearance of the flux*.

In conclusion, I believe that annihilation catastrophe may pretend to be not only one of the most dramatic manifestations of the reaction-diffusion interplay, but it is a *new type of catastrophe* wherein two explosive processes (Ω_{Hs} and Ω_{Ls}) are developing simultaneously, effectively "compensating" one another so that for an external observer (J) the explosion dynamics goes unnoticed up to the point of singularity, in the vicinity of which decompensation of explosions is manifested as a sudden singular jump.

This work was supported by the RFBR Grant No. 02-03-33122-a.

References: [1] For a review see E. Kotomin and V. Kuzovkov, *Modern Aspects of Diffusion Controlled Reactions: Cooperative Phenomena in Bimolecular Processes* (Elsevier, Amsterdam, 1996); [2] A.A. Ovchinnikov and Ya.B. Zeldovich, Chem. Phys. **28**,

215 (1978); [3] D. Toussaint and F. Wilczek, J. Chem. Phys. **78**, 2642 (1983); [4] L. Galfi and Z. Racz, Phys. Rev. A **38**, 3151 (1988); [5] E. Ben-Naim and S. Redner, J. Phys. A **25**, L575 (1992); [6] S. Cornell and M. Droz, Phys. Rev. Lett. **70**, 3824 (1993); [7] B.M. Shipilevsky, Phys. Rev. Lett. **73**, 201 (1994); [8] B.M. Shipilevsky, Phys. Rev. Lett. **82**, 4348 (1999); [9] B.M. Shipilevsky, J. Phys. A **30**, L471 (1997); [10] B.M. Shipilevsky (to be published); [11] L.P. Kadanoff, *Singularities and Blowups*, Phys. Today, 11 (1997).